




Care Home Infection Prevention and Control (IPC) Resource for Respiratory illness



Version 4.0
October 2025

Version History

Version	Date	Summary of changes
1.0	30 November 2023	Final
2.0	7 October 2024	Removal of 2m guidance for cohort to align with NIPCM amendments Reference to revised Transmission Based Precautions Definitions Updated Appendices to align to NIPCM/CHIPCM amendments Link to new NIPCM winter preparedness webpage.
3.0	10 October 2025	Amended wording for version control Updated link to winter preparedness page for 2025 Add sentence regarding transfer of infectious residents to other care facility
4.0	31 October 2025	Amended wording for version control Amended wording to reflect current Scottish immunisation schedule, and updated links Added sentence regarding transfer of infectious residents to other care facility

ARHAI Scotland are currently undertaking a Transmission-based Precautions (TBPs) definitions literature review and developing recommendations for practice. It is likely that 'droplet transmission' and 'airborne transmission' will be replaced with new definitions to describe respiratory transmission.

This will mean changes throughout the Care Home Infection Prevention and Control Manual (CH IPCM) to update the terminology including the addition of resources to support any guidance changes.

Using this resource online is advised as it ensures materials accessed will remain up to date.

All national Infection Prevention and Control (IPC) updates for the care home sector are communicated to stakeholders via the Care Home IPC Oversight and Advisory Group in addition to being uploaded to the [news section of the NIPCM.](#)

Contents

Section 1: Introduction	5
Section 2: Seasonal Preparation	6
Section 3: Infection Prevention and Control (IPC).....	7
3.1 Standard Infection Control Precautions (SICPs)	7
3.2 Transmission-based Precautions (TBPs).....	8
3.3 Hierarchy of Controls (HoC)	9
Section 4: Respiratory Infection Outbreaks	10
4.1 Case definitions	11
4.2 Actions to be taken in the event of a suspected or confirmed outbreak	11
Appendix 1: Seasonal IPC Checklist for RI	13
Appendix 2: RI IPC Outbreak Checklist	15
Appendix 3: RI Outbreak Care Monitoring Record IPC Outbreak Checklist	19
Appendix 4: RI Outbreak Case Review Monitoring Data Record	20

Section 1: Introduction

This care home respiratory illness (RI) infection prevention and control (IPC) resource has been developed by ARHAI Scotland to provide care home staff with key IPC information to support the management of suspected and confirmed cases of respiratory illness of an infectious nature amongst residents. This resource should be used alongside the [Care Home Infection Prevention and Control Manual \(CH IPCM\)](#).

Infectious respiratory illness can be caused by bacteria, viruses, and less commonly fungus. Symptoms, incubation periods, and the period of infectivity may vary between pathogens and individuals.

The most common causative pathogens are viral and include influenza (flu), Respiratory Syncytial Virus (RSV) and COVID-19.

Infectious Respiratory illness should be suspected in any individual with either a high temperature (fever) or sudden decline in mental or physical ability/functioning, in combination with any new onset of respiratory symptoms, for example cough and/or sore throat. Residents should be clinically assessed by a GP or via existing processes to exclude any differential diagnosis.

Understanding incubation periods and periods of infectivity will support resident assessment, resident placement, and enable the identification of residents who may have been exposed for ongoing symptom monitoring and early detection of new cases. Full details regarding pathogen specific symptoms, incubation periods and periods of infectivity are listed in the national [A-Z pathogens](#).

Section 2: Seasonal Preparation

Infectious respiratory illnesses can occur at any time throughout the year, although influenza and RSV are often a winter season virus. Staff should be continually aware and remain symptom vigilant for respiratory infections in the care home and the possibility of an outbreak.

[Winter vaccinations](#) are available for flu and COVID-19 and remain one of the best ways to protect individuals. The vaccines are offered annually and can protect the vulnerable from developing severe illness and from requiring a hospital admission's vaccinations.

Older adults and vulnerable groups are now also eligible for the RSV vaccine which has been added to the Scottish vaccine schedule. To support care home provider and staff preparedness, winter IPC communication reminders and corresponding support materials are issued annually by ARHAI Scotland and NHS Education for Scotland (NES). Once published these live links will be promoted and disseminated for local use to support seasonal preparation plans in care homes to help raise awareness and to support education and training for care home staff for all relevant communications, resources, including supportive education and training for care home staff.

Care homes providers and staff are advised to prepare for seasonal RI illness by reviewing and considering all the necessary actions in [Appendix 1](#) of this guidance to protect all staff and residents, especially those who are at highest risk of becoming seriously ill.

Section 3: Infection Prevention and Control (IPC)

Respiratory infections can be transmitted or spread in various ways. Staff should be familiar with infection transmission routes.

The care home designated person in charge should ensure that IPC precautions are adopted and implemented via ongoing IPC compliance monitoring.

There should also be local processes in place for preadmission assessment/ triage of infection risks for **all residents** prior to arrival (where possible) or carried out on arrival. It is important to be vigilant for all new admissions or returning residents for any symptoms of infection. As a reminder, it is also important to undertake the [general respiratory screening questions](#) before a resident is admitted to the care home.

To prevent the unintended introduction or spread of infection from the care home, where possible, advanced warning should be provided to the Scottish Ambulance Service, and other healthcare facilities (including hospitals) in advance of transfer, for resident(s) with known RI infection or symptoms suggestive of a probable infection.

3.1 Standard Infection Control Precautions (SICPs)

The basic IPC measures that should be used in the care home are called [Standard Infection Control Precautions](#) (SICPs). When applied, these reduce the risk of transmission of infectious agents from known and unknown sources of infection.

SICPs should be used by **all staff, in all care settings, at all times, for all residents**, whether infection is known to be present or not to ensure the safety of residents, staff and visitors in the care home.

It is essential that SICPs are applied continuously as residents living in care homes are more vulnerable, therefore increasing their risk of acquiring infections which may then be serious and potentially life threatening. By continuously applying SICPs, you will provide a safe environment and effective care.

If you suspect or are aware that a resident has an infection, then details should be confirmed for the correct IPC precautions to be put in place for the safety of the resident and others. Obtaining infection details may include appropriate clinical samples and/or screening to establish the causative organism which may be on advice from your local GP, Health Protection Team (HPT) and/or Infection Prevention and Control Team (IPCT) (depending on local processes).

If you suspect a resident has a RI you should:

- inform the person in charge immediately
- commence transmission-based precautions
- encourage and assist residents and those providing meaningful contact to practice good hand hygiene and respiratory hygiene
- continue to carry out hand hygiene as per [CH IPCM](#) at the [WHO 4 Moments](#) using hand rub unless they are visibly soiled, otherwise wash hands with soap and water
- use the appropriate personal protective equipment (PPE) when carrying out direct care as outlined in [Appendix 15](#)
- ensure timely collection of specimens for symptomatic residents which may include throat or perinasal swabs

For further information about SICPs see [Chapter 1 of the CH IPCM](#)

3.2 Transmission-based Precautions (TBPs)

When caring for a resident who presents with symptoms of a respiratory illness, not in keeping with their normal respiratory function and therefore has a suspected or known RI illness, SICPs won't be enough to stop infection spreading and you will need to use some extra IPC precautions. These extra precautions are called [Transmission-Based Precautions](#) (TBPs).

Clinical judgement and decisions should be made by staff to determine the necessary IPC precautions required (the local HPT and/or IPCT (depending on local processes) should be contacted for advice and support where required).

Clinical judgement and decisions should be based on the:

- suspected or known infectious agent
- transmission route of the infectious agent
- care setting and procedures undertaken
- severity of the illness caused

Prompt application of TBPs is essential to prevent further spread, and testing of symptomatic residents is required to confirm identification of the pathogen.

Symptomatic staff should refrain from duty.

Visitors should be supported with IPC and [PPE requirements](#).

Further information regarding TBPs see [Appendix 11](#) and [Appendix 15 of the NIPCM](#).

The local HPT or IPCT (depending on local processes) should be contacted for advice when required.

3.3 Hierarchy of Controls (HoC)

The [Hierarchy of Controls \(HoC\)](#) should be considered when applying SICPs and TBPs recognising that the most effective method of control (elimination) is employed first. This inherently results in safer control systems.

It is recognised that elimination of risk may be challenging within health and care settings due to the nature of the services provided. Where that is not possible, all other controls must be considered in sequence.

Personal protective equipment (PPE) is the **last** in the HoC and may be the only mitigating control when caring for a resident.

Further information on the HoC can found in the [CH IPCM](#) and [Appendix 17 of the NIPCM](#).

Section 4: Respiratory Infection Outbreaks

Definitions of infection outbreaks and incidents can be found in [Chapter 3 of the NIPCM](#).

For staff cases, a community source (i.e. out with the care home setting) should be considered and eliminated where possible.

Care homes for older adults are considered higher-risk settings for outbreak management purposes. As such, Health Protection Teams (HPTs) and/or Infection Control Teams (IPCTs) (depending on local processes) should be notified when a cluster of suspected or confirmed cases are identified.

This is due to:

- the population being older and likely to have more underlying health conditions. This puts them at greater risk of more severe illness
- opportunities for infections to spread quickly throughout the facility due to the communal nature of the setting

Further information can be found in Section 3.2 of [SHPN Management of public health incidents. Guidance on the Roles and Responsibilities of an NHS led Incident Management Team](#).

The local HPT and/or IPCT (depending on local processes) should be contacted for further advice or support where required.

Staff should refer to the agreed minimum [NHS Alert organism/condition list in Appendix 13 of the NIPCM](#) for respiratory pathogens that require further investigation and reporting.

4.1 Case definitions

Case definitions for assessment and reporting purposes will be defined by the supporting HPT/IPCT and/or the incident management team (IMT).

Examples are provided below.

- **Confirmed case.** Any resident or staff member who has or had respiratory symptoms of an infectious nature and has tested positive for a respiratory pathogen with epidemiological links to the care home setting.
- **Suspected case.** Any resident or staff member with respiratory symptoms assessed as being of an infectious nature who has not yet been laboratory confirmed that has epidemiological links to any confirmed case.

4.2 Actions to be taken in the event of a suspected or confirmed outbreak

The care home designated person in charge should ensure that:

- symptomatic residents are clinically assessed in line with local agreed processes
- IPC precautions are in place
- individual documented risk assessments are in place for all symptomatic residents who require to be isolated in their individual rooms. Risk assessments should consider individual resident health and wellbeing, their health and safety, and their ability to remain in their rooms with the door closed. The minimum time for isolation in line with the [A-Z of pathogens](#) from resolution of symptoms should be supported and reviewed daily
- external receiving staff are notified if the resident being transferred has had a suspected or confirmed RI pathogen within the previous 48 hours
- symptomatic staff do not report for duty

The local HPT and/or IPCT (depending on local processes) should be contacted for further advice or support where required in relation to outbreak management. This may include testing for symptomatic residents to identify the causative organism that will inform individual placement if cohorting is required and direct any required treatment or any specialist advice.

The RI outbreak checklist in [Appendix 2](#) is a support resource which may be used to support compliance with IPC measures.

[Appendix 3](#) and [Appendix 4](#) should also be used to support assessment and record keeping.

Appendix 1: Seasonal IPC Checklist for RI

Number	Actions to prepare	Y	N	Date
1	Has this resource and associated winter IPC preparedness been communicated to all staff?			
2	Are all staff aware of and have access to the CH IPCM ?			
3	Have all staff completed appropriate IPC training relevant to their roles?			
4	Have all residents and staff been encouraged (where appropriate) to uptake winter vaccinations and have been provided with details of NHS Inform information about the available seasonal vaccinations, including processes for vaccination booking information?			
5	Have all residents who are at highest risk of becoming seriously ill from respiratory pathogens been identified and assessed?			
6	Has PHS guidance for consent in care homes in Scotland been implemented and available in advance of vaccination schedules?			
7	Have resident seasonal vaccinations been scheduled and delivered as appropriate?			
8	Is PPE available in sufficient quantities and stored in a clean/dry area until required for use?			
9	Have all relevant staff been fit tested for respiratory protective equipment and trained to perform a fit check?			
10	Have hand washing posters as per the CH IPCM and WHO 4 moments posters been displayed in key areas throughout the care home?			
11	Is there liquid soap and disposable paper hand towels available in all public and communal toilets and in en-suite toilets?			
12	Are there sufficient quantities of hand rub and disposable tissues available in all public and communal areas and in resident's rooms?			
13	Are lidded foot operated waste bins in place and functioning throughout the facility?			

Number	Actions to prepare	Y	N	Date
14	Are facilities in place for segregation of infectious laundry and water-soluble laundry bags readily available?			
15	Are there sufficient quantities of required cleaning materials available?			
16	Are there processes in place to support symptomatic resident risk assessments?			
17	Are ARHAI Scotland Transmission Based Precautions posters for droplet precautions displayed in key areas (throughout the care home for the duration of the outbreak)?			
18	Are ' Catch it, Bin it, Kill it ' posters displayed in highly visible areas?			

Appendix 2: RI IPC Outbreak Checklist

The IPC checklist should be completed as soon as an outbreak is suspected and continued daily until declared over.

State pathogen(s) and date the outbreak was identified:

Transmission Based Precautions for Outbreak Management
SICPs should continue to be applied in all care settings, at all times, for all residents

Action	Date				
Resident Placement/Assessment of risk					
Residents who are confirmed, suspected and those at high risk of acquisition and adverse outcomes from infection (e.g. immunosuppressed) have been prioritised for single room accommodation with ensuite facilities					
Hierarchy of controls has been assessed and additional controls implemented where risk assessed and able to do so (e.g. window opening)					
Doors to isolation/cohort rooms/areas are closed and signage is clear. Individual resident risk assessments are documented in the resident notes for door closure in relation to resident safety, and health and well-being, minimum period specified and reviewed daily.					
Cohort areas are established if resident single rooms are unavailable and where there are multiple cases of the same infection.					
Contact tracing for exposed residents has been undertaken with symptom monitoring in place for early detection of new cases.					



Action	Date				
Personal Protective Equipment (PPE): gloves, disposable aprons, fluid resistant type II surgical mask (FRSM), eye/face protection and respiratory protection (RPE)					
Staff are wearing and applying correct use of PPE for direct care contact or when in the residents immediate care environment and PPE is changed between residents and/or following completion of a procedure or task – disposable apron, FRSM, eye/face protection. The use of gloves has been risk assessed.					
Staff are wearing and applying correct use of RPE as per the CH IPCM including the correct duration for post AGP fallow times - fluid repellent gown, fluid resistant FFP3 respirator, eye/face protection, and gloves.					
Staff are aware that where RPE (FFP3 masks) is required, they are to be doffed following exit from the care environment when the exposure risk has ended.					
Sufficient stocks of PPE including RPE are easily located and stored from risk of contamination.					
Symptomatic residents are encouraged to wear FRSMs where tolerated.					
Safe Management of Care Equipment					
Single-use items are in use.					
Dedicated reusable non-invasive care equipment is in use and decontaminated between use and prior to use on another resident using detergent followed by a disinfectant solution of 1000 parts per million available chlorine (ppm av cl) or a combined detergent/disinfectant solution at a dilution of 1000 ppm av cl.					
Fans are not in use or risk assessments are in place in relation to clinical need.					
Safe Management of the Care Environment					
All areas are free from non-essential items for resident care and equipment; exposed food stuffs have been removed.					

Action		Date				
At least daily cleaning and decontamination of the resident isolation room/cohort rooms/areas is in place using detergent followed by a disinfectant solution of 1000 ppm av cl or a combined detergent/disinfectant solution at a dilution of 1000 ppm av cl. Discuss with local HPT/ICT if items are unable to withstand chlorine releasing agents.						
Increased frequency of decontamination is incorporated into the environmental decontamination schedules for areas where there may be higher environmental contamination rates, for example "frequently touched" surfaces such as door/toilet handles, bedside tables, over bed tables and bed rails.						
Terminal decontamination is undertaken following resident transfer, discharge, or once the resident is no longer considered infectious.						
Information and Treatment						
Residents						
Residents are informed of all screening/investigation result(s) where appropriate and the next of kin informed where consent has been provided. (Documented in resident notes)						
Resident Information Leaflet has been provided and explained to next of kin where consent is in place. (Document in resident notes).						
Any appropriate antimicrobial/antiviral prophylaxis and treatment has been administered as prescribed.						
Symptomatic residents are supported to wear a FRSM where clinical safe and if tolerated.						
Staff						
Staff are kept informed of updates relating to the outbreak and necessary controls.						

Action	Date				
Staff considered at increased risk of acquisition (immunosuppression etc) have been risk assessed by Care Home Manager for resident care allocation.					
No symptomatic staff are on duty and the A-Z pathogen exclusion guidance is followed.					
Visitors					
All visitors are aware of the current situation.					
IPC support and advice has been offered to those providing direct care to a resident. FRSM and eye/face protection has been offered.					
Visitors have been informed if they develop symptoms they should stay at home and follow advice by NHS inform .					
Provide relatives and those providing meaningful contact with the Washing Clothes at Home Leaflet					
Staff support visitors with Scottish Government Care Home Visiting Policy					

(Refer to the [Care Home IPC Manual](#) for further information)

Appendix 3: RI Outbreak Care Monitoring Record IPC Outbreak Checklist

Case assessments.

Specify Respiratory Pathogens:

Residents:	Date						
Number of new probable residents today							
Total number of probable residents							
Number of new confirmed cases today							
Total number of new confirmed case today							
Total number of remaining symptomatic cases today							
Total number of residents giving cause for concern as a direct result of incident today							
Number of new residents who have died as a direct result of the respiratory pathogen and has been reported on any part of the death certificate							
Total number of resident deaths with respiratory pathogen reported on part 1 or 2 of the death certificate to date							

Staff:	Date:						
Number of new symptomatic staff today							
Total number of staff affected to date							

Appendix 4: RI Outbreak Case Review Monitoring Data Record

Complete for all residents with symptoms.

Specify Respiratory Pathogen:

Name of resident/staff member	CHI	Vaccinated (Y/N)	Antibiotic (Y or N)	Antiviral (Y/N/NA)	Specimen date (X)	Specimen result (+ or -)	Date of start and end of symptoms (Identify start and end date with a dot as below and link dates with a line)						
Example							19/9	20/9	21/9	22/9	23/9	24/9	25/9
Joe Bloggs	0112326214	N	N	N	20/9	+		X					
Jane Brown	0712325321	Y	N	Y									